
Fifth Edition

Electrocardiography

FOR HEALTHCARE PROFESSIONALS



Kathryn Booth, RN-BSN, RPT, RMA (AMT), CPhT, MS Thomas E. O'Brien, AS, CCT, CRAT, RMA



ELECTROCARDIOGRAPHY FOR HEALTHCARE PROFESSIONALS

Fifth Edition

Kathryn A. Booth, RN-BSN, RMA (AMT), RPT, CPhT, MS

Total Care Programming, Inc. Palm Coast, Florida

Thomas E. O'Brien, AS, CCT, CRAT, RMA, CCMA

Remington College Medical Assisting Program, Lead Instructor







ELECTROCARDIOGRAPHY FOR HEALTHCARE PROFESSIONALS, FIFTH EDITION

Published by McGraw-Hill Education, 2 Penn Plaza, New York, NY 10121. Copyright © 2019 by McGraw-Hill Education. All rights reserved. Printed in the United States of America. Previous editions © 2016, 2012. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written consent of McGraw-Hill Education, including, but not limited to, in any network or other electronic storage or transmission, or broadcast for distance learning.

Some ancillaries, including electronic and print components, may not be available to customers outside the United States.

This book is printed on acid-free paper.

1 2 3 4 5 6 7 8 9 0 LMN 21 20 19 18

ISBN 978-1-260-06477-3 MHID 1-260-06477-8

Executive Portfolio Manager: William R. Lawrensen

Senior Product Developer: Christine Scheid Executive Marketing Manager: Roxan Kinsey Content Project Manager: Ann Courtney Senior Buyer: Susan K. Culbertson

Design: David W. Hash

Content Licensing Specialists: Lori Hancock Cover Image: ©Sofiia Balitckaia/Shutterstock.com

Compositor: SPi Global

All credits appearing on page or at the end of the book are considered to be an extension of the copyright page.

Library of Congress Cataloging-in-Publication Data

Names: Booth, Kathryn A., 1957- author.

Title: Electrocardiography for healthcare professionals / Kathryn A. Booth, RN-BSN, RMA (AMT), RPT, CPhT, MS, Total Care Programming, Inc., Palm Coast, Florida, Thomas O'Brien, AS, CCT, CRAT, RMA, CCMA, Remington College, Medical Assisting Program, Lead Instructor.

Description: Fifth edition. | New York, NY : McGraw-Hill Education, [2019] |

Includes index.

Identifiers: LCCN 2017036542 | ISBN 9781260064773 (alk. paper)

Subjects: LCSH: Electrocardiography.

Classification: LCC RC683.5.E5 H256 2019 | DDC 616.1/207547—dc23 LC record available at

https://lccn.loc.gov/2017036542

The Internet addresses listed in the text were accurate at the time of publication. The inclusion of a website does not indicate an endorsement by the authors or McGraw-Hill Education, and McGraw-Hill Education does not guarantee the accuracy of the information presented at these sites.



Dedication To the individuals using the information in this book, you have chosen a worthwhile and rewarding profession. Thank you; your skills and services are truly needed. To my children and grandchildren who keep me young. Kathryn Booth I want to thank my beautiful wife, Michele, and our wonderful children, Thomas, Robert, and Kathryn. Without their love and support, I would have nothing. They inspire me every day to make a difference in people's lives. I also want to express my sincere thanks to the faculty, staff, and students of Remington College for their encouragement and guidance. Today's students are the difference makers of tomorrow! Thomas O'Brien

About the Authors

Kathryn A. Booth, RN-BSN, RMA (AMT), RPT, CPhT, MS, is a registered nurse (RN) with CPR and ACLS training as well as a master's degree in education and certifications in phlebotomy, pharmacy tech, and medical assisting. She is an owner, author, educator, and consultant for Total Care Programming. Inc. She has over 35 years in healthcare and education including nursing, electrocardiography, phlebotomy, and medical assisting. Her experience spans five states. As an educator, Kathy has been awarded the teacher of the year in three states where she taught various health educational courses including healthcare science, nursing, and medical assisting. She serves on the Cardiovascular Credentialing International Certified Rhythm Analysis Technician Item Writer and Reviewer's committee and the American Medical Technologists registered Phlebotomy Technician Examinations, Qualifications, and Standards committee. She is a past and current member of two educational advisory boards. Her larger goal is to develop up-to-date, dynamic healthcare educational materials to assist other educators as well as to promote the healthcare professions. In addition, Kathy enjoys presenting innovative new learning solutions for the changing healthcare and educational landscape to her fellow professionals nationwide.

Thomas E. Davidson-O'Brien, AS, CCT, CRAT, RMA, CCMA, is the lead instructor for the Medical Assisting program at Remington College, Fort Worth, Texas. Tom also works as an author and editor of Practical Clinical Skills (www.practicalclinicalskills.com) and EKG academy. He is a former member of the Board of Trustees and Exam Chair for the Certified Cardiographic Technician and Certified Rhythm Analysis Technician Registry Examinations working with Cardiovascular Credentialing International (CCI). Tom continues to volunteer his efforts as an active contributor to both credential exams as a member of the Item Writing team. His background includes over 35 years in healthcare including the U.S. Air Force and U.S. Army Medical Corps. Tom's medical career as an Air Force Independent Duty Medical Technician (IDMT) has taken him all over the United States and the world. He has several years' experience working in the Emergency Services and Critical Care arena (Cardiothoracic Surgery and Cardiac Cath Lab). He was awarded Master Instructor status by the U.S. Air Force in 1994 upon completion of his teaching practicum. He now has over 18 years of teaching experience; subjects include Emergency Medicine, Cardiovascular Nursing, Fundamentals of Nursing, Dysrhythmias, and 12-Lead ECG Interpretation. His current position provides challenges to meet the ever-changing needs of the medical community and to provide first-rate education to a diverse adult education population.

Brief Contents

Preface xi Acknowledgments xviii Electrocardiography 1 CHAPTER The Cardiovascular System 33 CHAPTER The Electrocardiograph 59 CHAPTER Performing an ECG 90 CHAPTER Rhythm Strip Interpretation and Sinus Rhythms CHAPTER Atrial Dysrhythmias 157 CHAPTER Junctional Dysrhythmias 176 CHAPTER Heart Block Dysrhythmias CHAPTER Ventricular Dysrhythmias CHAPTER Bundle Branch Block 245 CHAPTER Pacemaker Rhythms 257 CHAPTER Cardiac Stress Testing 279 CHAPTER Ambulatory Monitoring CHAPTER 14 Clinical Presentation and Management of the Cardiac CHAPTER Patient 335 Basic 12-Lead ECG Interpretation 365 CHAPTER





Cardiovascular Medications A-1 APPENDIX Standard and Isolation Precautions B-1 APPENDIX Medical Abbreviations, Acronyms, and Symbols C-1 APPENDIX **Anatomical Terms D-1** APPENDIX Identifying Components of ECG Strips E-1 APPENDIX Glossary G-1 Index I-1

Contents

		Preface xi
		Acknowledgments xviii
		Acknowledginetits xviii
CHAPTER		Electrocardiography 1
	$-\!$	
	'	1.1 The ECG and Its History 2
		1.2 Uses of an ECG 3
		1.3 Preparing for an ECG 10
		1.4 Infection Control 15
		1.5 Vital Signs 19
		Competency Checklists
		1-1 Measuring Pulse and Respirations 29
		1-2 Measuring Adult Blood Pressure 31
		The Cardiavaccular System 22
CHAPTER	_ \	The Cardiovascular System 33
	 ~\	
		2.1 Circulation and the ECG 34
		2.2 Anatomy of the Heart 34
		2.3 Principles of Circulation 39
		2.4 The Cardiac Cycle 41
		2.5 Conduction System of the Heart 43
		2.6 Electrical Stimulation and the ECG Waveform 46
CHAPTER	3	The Electrocardiograph 59
CHAPIER	1	hile Liectrocardiograph 33
	7 ~ 1	
		3.1 Producing the ECG Waveform 59
		3.2 ECG Machines 65
		3.3 ECG Controls 70
		3.4 Electrodes 74
		3.5 ECG Graph Paper 75
		3.6 Calculating Heart Rate 78
CHAPTER	1	Performing an ECG 90
CHAFIER	٨	Terrorining an Eoo 50
	"\	
		4.1 Preparation for the ECG Procedure 91
		4.2 Communicating with the Patient 92

	4.3 Safety 94
	4.4 Applying the Electrodes and Leads 96
	4.5 Operating the ECG Machine 101
	4.6 Checking the ECG Tracing 102
	4.7 Reporting ECG Results 106
	4.8 Equipment Maintenance 108
	4.9 Pediatric ECG 109
	4.10 Cardiac Monitoring 110
	4.11 Special Patient Considerations 111
	4.12 Handling Emergencies 115
	Competency Checklists
	4-1 Recording an Electrocardiogram 124
	4-2 Continuous Cardiac Monitoring 126
_	3
CHAPTER	Rhythm Strip Interpretation and
	Sinus Rhythms 128
	5.1 Rhythm Interpretation 128
	5.2 Identifying the Components of the Rhythm 129
	5.3 Rhythms Originating from the Sinus Node 136
	5.4 Sinus Bradycardia 139
	5.5 Sinus Tachycardia 141
	5.6 Sinus Dysrhythmia 142
	5.7 Sinus Arrest 144
	5.8 Sinus Exit Block 147
	5.6 Silius Exit Block 147
CHAPTER 6	Atrial Dysrhythmias 157
CHAITER	
	Y
	6.1 Introduction to Atrial Dysrhythmias 157
	6.2 Premature Atrial Complexes 158
	6.2 Premature Atrial Complexes 158
	6.2 Premature Atrial Complexes 1586.3 Wandering Atrial Pacemaker 160
	6.2 Premature Atrial Complexes6.3 Wandering Atrial Pacemaker6.4 Multifocal Atrial Tachycardia162
_	 6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163
CHAPTER 7	 6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165
CHAPTER 7	 6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165
CHAPTER 7	6.2 Premature Atrial Complexes 6.3 Wandering Atrial Pacemaker 6.4 Multifocal Atrial Tachycardia 6.5 Atrial Flutter 6.6 Atrial Fibrillation 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176
CHAPTER 7	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176
CHAPTER 7	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176 7.2 Premature Junctional Complex (PJC) 178
CHAPTER 7	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176
CHAPTER 7	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176 7.2 Premature Junctional Complex (PJC) 178
CHAPTER 7	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176 7.2 Premature Junctional Complex (PJC) 178 7.3 Junctional Escape Rhythm 179
CHAPTER	6.2 Premature Atrial Complexes 158 6.3 Wandering Atrial Pacemaker 160 6.4 Multifocal Atrial Tachycardia 162 6.5 Atrial Flutter 163 6.6 Atrial Fibrillation 165 Junctional Dysrhythmias 176 7.1 Introduction to Junctional Dysrhythmias 176 7.2 Premature Junctional Complex (PJC) 178 7.3 Junctional Escape Rhythm 179 7.4 Accelerated Junctional Rhythm 181

CHAPTER 8	Heart Block Dysrhythmias 196
	 8.1 Introduction to Heart Block Dysrhythmias 196 8.2 First Degree Atrioventricular (AV) Block 196 8.3 Second Degree Atrioventricular (AV) Block, Type I (Mobitz I or Wenckebach) 198 8.4 Second Degree Atrioventricular (AV) Block, Type II (Mobitz II) 201 8.5 Third Degree Atrioventricular (AV) Block (Complete) 204
CHAPTER 9	Ventricular Dysrhythmias 216
	9.1 Introduction to Ventricular Dysrhythmias 216 9.2 Premature Ventricular Complexes (PVCs) 217 9.3 Agonal Rhythm 221 9.4 Idioventricular Rhythm 223 9.5 Accelerated Idioventricular Rhythm 224 9.6 Ventricular Tachycardia 226 9.7 Ventricular Fibrillation 228 9.8 Asystole 231
CHAPTER 10	Bundle Branch Block 245
44	10.1 Introduction to Bundle Branch Block Dysrhythmias 245 10.2 LBBB vs. RBBB 249
CHAPTER	Pacemaker Rhythms 257
	11.1 Introduction to Pacemakers 257 11.2 Evaluating Pacemaker Function 261 11.3 Pacemaker Complications Relative to the ECG Tracing 266 Competency Checklist 11-1 Pacemaker Evaluation 277
CHAPTER 12	Cardiac Stress Testing 279
	12.1 Cardiac Stress Testing 279 12.2 Why Is Exercise Electrocardiography Used? 281 12.3 Other Types of Cardiac Stress Testing 282 12.4 Preparing the Patient for Exercise Electrocardiography 285 12.5 Providing Safety 288 12.6 Performing Exercise Electrocardiography 291



ix

	12.7 Common Protocols 293
	12.8 After Exercise Electrocardiography 296
	Competency Checklist
	12-1 Assisting with Exercise Electrocardiography (Stress Testing) 304
43	
CHAPTER	Ambulatory Monitoring 308
	M
	43.4 Ambulatawa Manitavina 200
	13.1 Ambulatory Monitoring 308
	13.2 How Is Ambulatory Monitoring Used? 310
	13.3 Types of Ambulatory Monitoring 311
	13.4 Educating the Patient 316
	13.5 Preparing the Patient 319
	13.6 Applying an Ambulatory Monitor 320
	13.7 Removing an Ambulatory Monitor and Reporting Results 324
	Competency Checklist
	13-1 Applying and Removing an Ambulatory (Holter) Monitor 331
CHAPTER 14	Clinical Presentation and Management of the Cardiac
CHAPIER	
	Patient 335
	14.1 Coronary Arteries 336
	14.2 Cardiac Symptoms 338
	14.3 Atypical Patient Presentation 340
	14.4 Acute Coronary Syndrome 342
	14.5 Heart Failure 344
	14.6 Cardiac Patient Assessment and Immediate Treatment 347
	14.7 Treatment Modalities for the Cardiac Patient 352
	Competency Checklist
	14-1 Immediate Care for a Cardiac Patient 363
45	
CHAPTER	Basic 12-Lead ECG Interpretation 365
	·
	_
	15.4 Left Ventricular Hypertrophy 377
	Appendix A Cardiovascular Medications A-1
	Appendix B Standard and Isolation Precautions B-1
	Appendix C Medical Abbreviations, Acronyms, and Symbols C-1
	Appendix D Anatomical Terms D-1
	Appendix E Identifying Components of ECG Strips E-1
	Glossary G-1
	Index I-1

Preface

Healthcare is an ever-changing and growing field that needs well-trained individuals who can adapt to change. Flexibility is key to obtaining, maintaining, and improving a career in electrocardiography. Obtaining ECG training and certification, whether it be in addition to your current career or as your career, will make you employable or a more-valued employee. This fifth edition of *Electrocardiography for Healthcare Professionals* will prepare users for a national ECG certification examination, but most importantly, it provides comprehensive training and practice for individuals in the field of electrocardiography.

The fact that you are currently reading this book means that you are willing to acquire new skills or improve the skills you already possess. This willingness translates into your enhanced value, job security, marketability, and mobility. Once you complete this program, taking a certification examination is a great next step for advancing your career.

This fifth edition of *Electrocardiography for Healthcare Profes- sionals* can be used in a classroom as well as for distance learning. Checkpoint Questions and Connect exercises correlated to the Learning Outcomes
make the learning process interactive and promote increased comprehension.
The variety of materials included with the program provides for multiple
learning styles and ensured success.

Text Organization

The text is divided into 15 chapters:

- Chapter 1 Electrocardiography includes introductory information about the field as well as legal, ethical, communication, and patient education information. In addition, basic vital signs and troubleshooting are addressed. Competency checklists are included to practice checking vital signs.
- Chapter 2 The Cardiovascular System provides a complete introduction and review of the heart and its electrical system. The information focuses on what you need to know to understand and perform an ECG. Specific topics include anatomy of the heart, principles of circulation, cardiac cycle, conduction system and electrical stimulation, and the ECG waveform.
- Chapter 3 The Electrocardiograph creates a basic understanding of the ECG, including producing the ECG waveform, the ECG machine, electrodes, and ECG graph paper.
- Chapter 4 Performing an ECG describes the procedure for performing an ECG in a simple step-by-step fashion. Each part of the procedure is explained in detail, taking into consideration the latest guidelines. The chapter is divided into the following topics: preparation, communication, safety, anatomical landmarks, applying the electrodes and leads, operating the ECG machine, checking the tracing, reporting results, and equipment maintenance. Extra sections are included regarding pediatric

- ECG, cardiac monitoring, special patient circumstances, and emergencies. Competency checklists are included to practice performing both an ECG and continuous monitoring.
- Chapter 5 Rhythm Strip Interpretation and Sinus Rhythms introduces
 the five-step criteria for classification approach to rhythm interpretation
 that will be utilized throughout Chapters 5 through 11. With updated, realistic rhythm strip figures, explanations, and Checkpoint Questions, the
 user learns to interpret the sinus rhythms, including criteria for classification, how the patient may be affected, basic patient care, and treatment.
- Chapter 6 Atrial Dysrhythmias provides an introduction to and interpretation of the atrial dysrhythmias, including criteria for classification, how the patient may be affected, basic patient care, and treatment.
- Chapter 7 Junctional Dysrhythmias provides an introduction to and interpretation of the junctional dysrhythmias, including criteria for classification, how the patient may be affected, basic patient care, and treatment.
- Chapter 8 Heart Block Dysrhythmias provides an introduction to and interpretation of the heart block dysrhythmias, including criteria for classification, how the patient may be affected, basic patient care, and treatment.
- Chapter 9 Ventricular Dysrhythmias provides an introduction to and interpretation of the ventricular dysrhythmias, including criteria for classification, how the patient may be affected, basic patient care, and treatment.
- Chapter 10 Bundle Branch Block provides an introduction to bundle branch block dysrhythmias, including criteria for classification, how the patient may be affected, basic patient care, and treatment. The chapter also discusses how to differentiate between left and right bundle branch block.
- Chapter 11 Pacemaker Rhythms provides an introduction to pacemaker rhythms, evaluation of pacemaker function, and complications related to the ECG tracing.
- Chapter 12 Cardiac Stress Testing provides the information necessary to assist with the exercise electrocardiography procedure. The competency checklist provides the step-by-step procedure for practice and developing proficiency at the skill.
- Chapter 13 Ambulatory Monitoring includes the latest information about various types of ambulatory monitors and includes what you need to know to apply and remove a monitor. A procedure checklist is also provided for this skill.
- Chapter 14 Clinical Presentation and Management of the Cardiac Patient expands on the anatomy of the coronary arteries and relates them to typical and atypical cardiac symptoms. STEMI, non-STEMI, and heart failure are introduced. The chapter includes a section about sudden cardiac death as compared to myocardial infarction and finishes with assessment, immediate care, and continued treatment of the cardiac patient.
- Chapter 15 Basic 12-Lead ECG Interpretation provides an introduction to 12-lead ECG interpretation. It includes anatomical views of the coronary arteries and correlates the arteries with the leads and views obtained on a 12-lead ECG. It also identifies the morphologic changes in the tracing that occur as a result of ischemia, injury, and infarction. Axis deviation and left ventricular hypertrophy round out the chapter concepts. The last section helps users put all of these concepts together for 12-lead interpretation.

These chapters can be utilized in various careers and training programs. Following are some suggested examples:

- Telemetry technicians (Chapters 1–12, depending on requirements)
- EKG/ECG technicians (the entire book, depending on requirements)
- Medical assistants (the entire book, depending on where they work)
- Cardiovascular technicians working in any number of specialty clinics, such as cardiology or internal medicine (the entire book)
- Remote monitoring facilities personnel (Chapters 1–10, 13–15)
- Emergency medical technicians (Chapters 2, 5–10, 15, possibly more depending on where they work)
- Paramedics (Chapters 2–15)
- Nursing, especially for cross-training or specialty training (Chapters 2–15)
- Patient care tech or nursing assistant (Chapters 2–4, 13, perhaps more depending on job requirements)
- Polysomnography technologist (Chapters 2–10)
- Echocardiography technologist (Chapters 2, 5–12)
- Cardiac cath lab technologist (Chapters 2–11, 15)

New to the Fifth Edition

- Many revised figures, as well as new figures, tracings, and tables for improved clarity and up-to-date information that also provides additional student practice.
- Consolidated information on standard and isolation precautions.
- Expanded and reorganized information about pacemakers and bundle branch block in two separate chapters.
- Added and updated content about the following topics: sinus exit block, tilt test, telemetry (including mobile cardiac outpatient telemetry [MCOT]), cardiac stents, and left ventricular assist devices (LVAD).
- Added several competency checklists for student practice, including checklists for measuring pulse and respiration, measuring adult blood pressure, pacemaker evaluation, and immediate care for cardiac patients.

Features of the Text

- Key Terms and Glossary: Key terms are identified at the beginning of each chapter. These terms are in **bold**, **color** type within the chapter and are defined both in the chapter and in the glossary at the end of the book.
- Checkpoint Questions: At the end of each main heading in the chapter are short-answer Checkpoint Questions. Answer these questions to make sure you have learned the basic concepts presented.
- Think It Through: The Think It Through feature identifies problems
 and situations that may arise when you are caring for patients or performing a procedure. At the end of this feature, you are asked a question
 to answer in your own words.
- Safety & Infection Control: You are responsible for providing safe care and preventing the spread of infection. This feature presents tips and techniques to help you practice these important skills relative to electrocardiography.

- Communicate and Connect: Patient interaction and education and intrateam communication are integral parts of healthcare. As part of your daily duties, you must communicate effectively, both orally and in writing, and you must provide patient education. Use this feature to learn ways to perform these tasks.
- Law & Ethics: When working in healthcare, you must be conscious of the regulations of HIPAA (Health Insurance Portability and Accountability Act) and understand your legal responsibilities and the implications of your actions. You must perform duties within established ethical practices. This feature helps you gain insight into how HIPAA, law, and ethics relate to the performance of your duties.
- Real ECG Tracings: Actual ECG tracings, or rhythm strips, have been
 provided for easy viewing and to make the task of learning the various
 dysrhythmias easier and more realistic. Use of these ECG rhythm strips
 for activities and exercises throughout the program improves comprehension and accommodates visual learners.
- Competency Checklists: Checklists at the end of some chapters (as appropriate) help you practice your skills in performing key procedures.
- Chapter Summary: Once you have completed each chapter, take time
 to read and review the summary table. It has been correlated to key concepts and learning outcomes within each chapter and includes handy
 page number references.
- **Chapter Review:** Complete the chapter review questions, which are presented in a variety of formats. These questions help you understand the content presented in each chapter. Chapters 4, 11, and 12 also include Procedure Checklists for you to use to practice and apply your knowledge.

Resources

Additional Instructor Resources

- **Instructor's Manual** with course overview, lesson plans, answers to CheckPoint and End-of-Chapter Review questions, competency correlations, sample syllabi, and more.
- PowerPoint Presentations for each chapter, containing teaching notes
 correlated to learning outcomes. Each presentation seeks to reinforce
 key concepts and provide an additional visual aid for students.
- **Test Bank** and answer key for use in class assessment. The comprehensive test bank includes a variety of question types, with each question linked directly to a learning outcome from the text. Questions are also tagged with relevant topic, Bloom's Taxonomy level, difficulty level, and competencies, where applicable. The test bank is available in Connect, and Word and EZ Test versions are also available.
- Conversion Guide with a chapter-by-chapter breakdown of how the content has been revised between editions. The guide is helpful if you are currently using **Electrocardiography for Healthcare Professionals** and moving to the new edition, or if you are a first-time adopter.
- **Instructor Asset Map** to help you find the teaching material you need. These online chapter tables are organized by Learning Outcomes and allow you to find instructor notes, PowerPoint slides, and even test bank suggestions with ease! The Asset Map is a completely integrated tool designed to help you plan and instruct your courses efficiently and comprehensively. It labels and organizes course material for use in a multitude of learning applications.

All of these helpful materials can be found within your Connect course within the Library, under the Instructor Resources.




McGraw-Hill Connect® is a highly reliable, easy-touse homework and learning management solution that utilizes learning science and award-winning adaptive tools to improve student results.

Homework and Adaptive Learning

- Connect's assignments help students contextualize what they've learned through application, so they can better understand the material and think critically.
- Connect will create a personalized study path customized to individual SmartBook®.
- SmartBook helps students study more efficiently by delivering an interactive reading experience through adaptive highlighting and review.

Over **7 billion questions** have been answered, making McGraw-Hill Education products more intelligent, reliable, and precise.

Connect's Impact on Retention Rates, Pass Rates, and Average Exam Scores



Using **Connect** improves retention rates by **19.8%**, passing rates by **12.7%**, and exam scores by **9.1%**.

Quality Content and Learning Resources

- Connect content is authored by the world's best subject matter experts, and is available to your class through a simple and intuitive interface.
- The Connect eBook makes it easy for students to access their reading material on smartphones and tablets. They can study on the go and don't need Internet access to use the eBook as a reference, with full functionality.
- Multimedia content such as videos, simulations, and games drive student engagement and critical thinking skills.

73% of instructors who use **Connect** require it; instructor satisfaction **increases** by 28% when **Connect** is required.



@McGraw-Hill Education

Robust Analytics and Reporting

- Connect Insight® generates easy-to-read reports on individual students, the class as a whole, and on specific assignments.
- The Connect Insight dashboard delivers data on performance, study behavior, and effort. Instructors can quickly identify students who struggle and focus on material that the class has yet to master.
- Connect automatically grades assignments and quizzes, providing easy-to-read reports on individual and class performance.



©Hero Images/Getty Images



Impact on Final Course Grade Distribution

without Connect

22.9% A

31.0%

27.4% B

34.3%

22.9% C

18.7%

11.5% D

6.1%

15.4% F

9.9%

More students earn **As** and **Bs** when they use McGraw-Hill Education **Connect**.

Trusted Service and Support

- Connect integrates with your LMS to provide single sign-on and automatic syncing
 of grades. Integration with Blackboard®, D2L®, and Canvas also provides automatic
 syncing of the course calendar and assignment-level linking.
- Connect offers comprehensive service, support, and training throughout every phase of your implementation.
- If you're looking for some guidance on how to use Connect, or want to learn tips and tricks from super users, you can find tutorials as you work. Our Digital Faculty Consultants and Student Ambassadors offer insight into how to achieve the results you want with Connect.

Acknowledgments

Authors

Kathryn Booth: Thanks to all the reviewers who have spent time helping to make sure this fifth edition is up-to-date. In addition, I would like to acknowledge McGraw-Hill for supporting this book into its fifth edition and Jody James for being my right hand through the process.

Additionally, I would like to acknowledge Patricia Dei Tos and the members of the Inova Health system, who help to create and support the original development of this textbook. I would also like to thank my co-author, Thomas O'Brien, for his continued dedication and timely support for the project.

Thomas E. Davidson-O'Brien: I would like to acknowledge Mr. David Rubin, president & CEO of Aerotel Medical Systems (1998) Ltd., 5 Hazoref St., Holon 58856, Israel. I would like to express my sincere appreciation to a pair of former students and Central Florida Institute graduates: Rebecca Walton, CCT, for her contribution of Interpret-Tips and Jamie Merritt, CCT, for "bunny branch block." I would also like to give a special thank you to the staff members of the Non-Invasive Cardiology Departments at the Pepin Heart Hospital, Morton Plant Hospital, All Children's Hospital, and Palms of Pasadena Hospital for their inputs and generous donation of their time and expertise.

Additionally I would like to thank my co-workers and the leadership at Remington College, especially my Program Director Mrs. Luz Rangel and esteemed colleague Mr. Davy Hobson. A special thank you to Mr. John Michael Maloney, RCIS, and Mrs. Kathy Hellums, RCS, for their contributions of support and expertise without reservation.

Finally, a very special thank you to my son Rob for his hours devoted to scanning many of the cardiac rhythms in this text.

Consultants

Cynthia T. Vincent, MMS, PA-C
Alderson Broaddus College,
Philippi, WV
Jennifer Childers, MS PAC
Alderson Broaddus College,
Philippi, WV
Susan Hurley Findley, RN, MSN
Houston, TX
Lynn M. Egler, RMA, AHI, CPhT
St. Clair Shores, MI
Kimberly Speiring, MA
St. Clair Shores, MI

Fifth Edition Reviewers

Bernard Beckerman, MD York College, CUNY

Loretta Beneby

Legacy Education LLC, dba High Desert Medical College, CA

Stanley Brown

Mississippi State University

Jo Ann Carroll, RT(R), CPhT, NCPT, CET

American College of Education, IN

Alia Carter

Atlanta Primary Care

Nancy Castaneda

Arizona Medical Training Institute

Sue Coleman, LPN, AS

American National University, VA

Jamie Cuda, AAS, BS, MS, Ed.D.

Mohawk Valley Community College, NY

Theresa Errante-Parrino, CMA (AAMA), PMD, CPhT, EdD

Indian River State College, FL

Nancy Feulner, MS Ed MT(ASCP)

MS Ed

College of DuPage, IL

Jade Hernandez, MSN, APRN, AGACNP-BC

University of Louisiana-Lafayette

Alicia Kelley, MS PA-C

Massachusetts College of Pharmacy and Health Sciences

Lori King, MA

Great Lakes Institute of Technology, PA

Debra LeHew, CMA, AAMA

Anoka Technical College, MN

Marta Lopez, MD, LM, CPM, RMA,

Miami Dade College, FL

Monica Maldari, ACSM credentialed CEP & RCEP

Fitchburg State University, MA

Mary Marks, APRN

Mitchell Community College, NC

Carole Mullins, MPA Health

Administration

Southwestern Michigan College

Ashita Patel, BSc in Medical Laboratory Science

Wake Tech Community College, NC

Greg Russell

Front Range Community College, CO

Paul Visich, PhD, MPH

University of New England, ME

Danielle Wilken, MT (ASCP)

Goodwin College, CT

Previous Editions Reviewers

Emil P. Asdurian, MD Bramson ORT College Forest Hills, NY

Vanessa J. Austin, RMA, CAHI Clarian Health Sciences Center,

Medical Assistant Indianapolis, IN

Rhonda J. Beck, NREMT-P

Central Georgia Technical College

Macon, GA

Nia Bullock, PhD

Miller-Motte Technical College

Cary, NC

Jesse A. Coale, PA-C

Philadelphia University Philadelphia, PA Stephen Coleman

Central Florida Institute Palm Harbor, FL

Harvey Conner, AS, REMT-P

Oklahoma City Community College Oklahoma City, OK

Barbara S. Desch, LVN, CPC, AHI

 $San\ Joaquin\ Valley\ College\ Inc.$

Visalia/Hanford Campus

Visalia, CA

Melissa L. Dulaney

MedVance Institute of Baton Rouge

Baton Rouge, LA

Mary Patricia English

Howard Community College

Columbia, MD

Michael Fisher, Program Director

Greenville Technical College

Greenville, SC

Donna L. Folmar

Belmont Technical College

St. Clairsville, OH

Anne Fox Maric College Carson, CA

James R. Fry, MS, PA-C

Marietta College Marietta, OH

Michael Gallucci, MS, PT

Assistant Professor of Practice, Program in Physical Therapy

School of Public Health, New York

Medical College New York, NY

Jonathan I. Greenwald

Arapahoe Community College

Littleton, CO Grace Haines National College Dayton, OH

Linda Karp

Atlantic Cape Community College

Mays Landing, NJ

Deborah Kufs, MS, BSN, CEN, EMT-P

Hudson Valley Community College Troy, NY

Susie Laughter, BSN, RN

Cambridge Institute of Allied Health

Longwood, FL Elizabeth Laurenz National College Columbus, OH

Sheri A. Melton, PhD West Chester University

West Chester, PA

Stephen J. Nardozzi

Westchester Community College

Valhalla, NY

David James Newton, NREMT-P

Dalton State College

Dalton, GA

R. Keith Owens

AB-Tech Community College

Asheville, NC

Douglas A. Paris, BS, NREMT-P

Greenville Technical College

Department of Emergency Medical

Technology Greenville, SC

David Rice, AA, BA, MA

Career College of Northern Nevada

Reno, NV

Dana M. Roessler, RN, BSN Southeastern Technical College

Glennville, GA

Wayne A. Rummings, Sr. *Lenoir Community College*

Kingston, NC

David Lee Sessoms, Jr., MEd, CMA

Miller-Motte Technical College

Cary, NC

Mark A. Simpson, NREMT-P, RN,

CCEMTP
Director of EMS

 $Northwest ext{-}Shoals\ Community$

College

Muscle Shoals, AL

Linda M. Thompson, MS, RRT

 $Madison\ Area\ Technical\ College$

Madison, WI

Dyan Whitlow Underhill, MHA, BS

Miller-Motte Technical College

Cary, NC

Eddy van Hunnik, PhD

Gibbs College Boston

Boston, MA

Suzanne Wambold, PhD, RN RDCS,

FASE

The University of Toledo

Toledo, OH
Danny Webb
Milan Institute
Visalia, CA

Danielle Schortzmann Wilken

Goodwin College East Hartford, CT Stacey F. Wilson, MT/PBT (ASCP), CMA

Cabarrus College of Health Sciences

Concord, NC

Fran Wojculewicz, RN, BSN, MS

Maricopa Community College

Glendale, AZ

Roger G. Wootten

 $Northeast\ Alabama\ Community$

College Rainsville, AL

First Edition

Civita Allard

Mohawk Valley Community College

Utica, NY

Vicki Barclay

West Kentucky Technical College

Paducah, KY

Nina Beaman

Bryant and Stratton College

Richmond, VA

Cheryl Bell

Sanz School

Washington, DC

Stephanie Bernard, BA, NCMA

Sanford-Brown Institute

Jacksonville, FL

Gaule Carr, CPFT, RRT, MS

Illinois Central College

East Peoria, IL

Cyndi Caviness, CRT, CMA (AAMA), AHI

Montgomery Community College

Troy, NC

Harvey Conner, NRP, AHA

Oklahoma City Community College

Oklahoma City, OK

Mary Hewett, BSEMS, MEd, NREMTP

University of New Mexico

Albuquerque, NM

Charles Hill, Paramedic

North Georgia Technical College

Clarksville, GA

Cynthia Hill, MBA, CPT, CEHRS, CMA

NewBridge Cleveland Center for Arts

& Technology

Cleveland, OH

Scott Jones, BS, MBA, EMT-P

Victor Valley College

Apple Valley, CA

Konnie King Briggs, CCT, CCI; PBT, ASCP; CPCI, ACA

Houston Community College

Houston, TX

Myrna Lanier

Tulsa Community College

Tulsa, OK

Joyce Lockwood, NREMTP

Prince George's Community College

Largo, MD

David Martinez, Medical Assistant/

EMT

Vista College

Richardson, TX

John McBryde, Nationally Registered Paramedic

East Mississippi Community College

Mayhew, MS

Cheryl McQuay, CPT, CMA, CPI,

CEKG

Star Career Academy

Brick, NJ

Sheri Melton, PhD, ACSM

Certified Exercise Specialist,

ACSM Certified Health Fitness

Specialist

West Chester University

West Chester, PA

Bharat Mody, MD

Star Career Academy

Clifton, NJ

Nicole Palmieri, RN, AHI

 $Advantage\ Career\ Institute$

Eatontown, NJ

Lucy Della Rosa

Concorde Career Institute

Lauderdale Lakes, FL

Debra Shafer Suzanne Wambold, RN, PhD Blair College The University of Toledo Colorado Springs, CO $Toledo,\ OH$ Stephen Smith, MPA, RT, RRT Andrew Wood, MS, NREMT-P Stony Brook University Emergency Medical Training Stony Brook, NY Professionals, LLC $Lexington,\,KY$ Scott Tomek, Paramedic Century College White Bear Lake, MN

Electrocardiography

Learning Outcomes	1.1 Describe the history and the	e importance of the ECG.			
	1.2 Identify the uses of an ECG and opportunities for an electrocardiographer.				
	1.3 Troubleshoot legal, ethical, patient education, and communication issues related to the ECG.1.4 Perform infection control measures required for the ECG.				
	Key Terms	auscultated blood pressure	event monitor		
	automatic external	healthcare providers			
	defibrillator (AED)	hypertension			
	cardiac output	hypotension			
	cardiopulmonary	isolation precautions			
	resuscitation (CPR)	law			
	cardiovascular disease (CVD)	libel			
	cardiovascular technologist	medical professional liability			
	Code Blue	myocardial infarction (MI)			
	coronary artery disease (CAD)	(heart attack)			
	defibrillator	personal protective equipment			
	diastolic blood pressure	(PPE)			
	dysrhythmia	slander			
	ECG monitor technician	standard precautions			
	electrocardiogram (ECG)	stat			
	electrocardiograph	systolic blood pressure			
	electrocardiograph (ECG)	telemedicine			
	technician	tilt table test			
	ethics	vital signs			

(CVD) Disease related to the heart and blood vessels (veins and arteries).

coronary artery disease

(CAD) Narrowing of the arteries of the heart, causing a reduction of blood flow.

myocardial infarction (MI) (heart attack) A blockage of one or more of the coronary arteries causing lack of oxygen to the heart and death of the muscle tissue.

electrocardiograph An instrument used to record the electrical activity of the heart.

electrocardiogram (ECG) A tracing of the heart's electrical activity recorded by an electrocardiograph.

1.1 The ECG and Its History


According to the CDC, the leading cause of death in the United States every year since 1921 is **cardiovascular disease (CVD)**, or a disease of the heart and blood vessels. Approximately 610,000 Americans die every year because of **coronary artery disease (CAD)**, which is narrowing of the arteries of the heart, causing a reduction of blood flow. Unbelievably, one out of every three American adults has some form of CAD. You may know someone who has a heart condition. Maybe someone you know has had a **myocardial infarction (MI)** or **heart attack**.

An **electrocardiograph** is an instrument that allows the heart's electrical activity to be recorded and studied. It is used to produce an electrical (electro) tracing (graph) of the heart (cardio). This representative tracing of the electricity as it moves through the heart is known as an **electrocardiogram (ECG)**.

In 1887, an English physician, Dr. Augustus D. Waller (1856–1922), showed that electrical currents are produced during the beating of the human heart and can be recorded. Willem Einthoven (1860–1927) invented the first electrocardiograph, sometimes referred to as EKG (electro), Kardio (Greek for heart) gram (recording). This invention resulted in a Nobel Prize in Physiology or Medicine in 1924. Advancements in this technology have brought about today's modern ECG machines (see Figure 1-1). Technology continues to improve the availability and speed of computer interpretation and quickly communicates this information to a healthcare professional. Digital communication allows healthcare professionals to monitor patients from locations miles away.

Figure 1-1 Today's 12-lead ECG machine is attached to the patient's chest, arms, and legs using electrodes and lead wires. It records a tracing of the electrical activity of the heart.

©James Davies/Alamy Stock Photo



Copyright © 2019 by McGraw-Hill Education

Performing the actual ECG procedure is not difficult; however, it must be performed competently. The tracing of the electrical current of the heart must be accurate because it is used to make decisions about a patient's care. An inaccurate tracing could result in a wrong decision about the patient's medication or treatment. These decisions could result in a negative outcome for the patient.



- 1. What is the leading cause of death in the United States?
- **2.** Who is credited with determining that the human heart produces electrical currents that can be recorded?


healthcare provider The scope of practice of each healthcare provider will determine the extent of the interpretation and treatment of each of the cardiac dysrhythmias or conditions. Each specific scope is determined by the licensure of each state. For example: Prescribing a medication is the responsibility of the physician. But some state practices allow nurse practitioners and physician assistants to prescribe medication under the guidance of the physician.

1.2 Uses of an ECG

Healthcare providers study the ECG tracing to determine many things about the patient's heart. They look for abnormalities in the recording and changes from earlier recordings when available. The American Heart Association (AHA) recommends that individuals over the age of 40 have an ECG done annually as part of a complete physical. This baseline tracing assists the physician in diagnosing abnormalities of the heart. A sample of a normal tracing is shown in Figure 1-2.

Electrocardiography can be performed in a number of healthcare settings. The type of ECG tracing produced depends on the setting and the type of ECG machine used.

Figure 1-2 A normal ECG tracing is a horizontal line with upward and downward waves or deflections that indicate electrical activity within the heart.



Copyright © 2019 by McGraw-Hill Education

Code Blue An alert that notifies healthcare providers that a patient is unresponsive and needs assistance immediately.

stat Immediately.

In the Hospital (Acute Care)

The 12-lead ECG is one of the most commonly used tools to assess the heart in the hospital setting. A 12-lead ECG provides a tracing of the electrical activity in the patient's heart. In the hospital, routine ECGs are frequently done before surgery. A 12-lead ECG also may be performed during an emergency frequently called a **Code Blue**. An emergency ECG may be required **stat**, or immediately. These are done for many reasons including when a patient experiences chest pain, shortness of breath, dizziness, syncope, altered mental status, or has a change in cardiac rhythm.

Think It Through



Remain Calm

It is essential that you remain calm when recording a stat ECG. Remaining calm is necessary to avoid stress to the patient and to reduce confusion during the emergency.

What would be an appropriate way to tell a patient you are doing a stat ECG?

Another use of the ECG tracing in the hospital is during continuous cardiac monitoring. The purpose of continuous monitoring is to observe the pattern of the electrical activity of the patient's heart over time. During continuous monitoring, sensors, also known as *electrodes*, are attached to the patient's chest. Wires are connected to the sensors and a small transmitter-like box or connected directly to a monitor displaying the patient's heart rhythm. Patients on continuous monitoring are usually in an intensive care unit (ICU), coronary care unit or cardiac care unit (CCU), surgical intensive care unit (SICU), inpatient telemetry unit, or emergency department (ED). Continuous monitoring is also done routinely during surgery.

Continuous monitoring done in a hospital with a small transmitter box connected to the patient using electrodes and lead wires is known as *telemetry monitoring*. The monitor is usually housed in a case and attached to the patient so he or she can move about. The ECG tracing is transmitted to a central location for evaluation. When several patients are on a telemetry unit, the tracings of all the patients are recorded on multiple monitors at the nursing or patient care station.

Doctors' Offices and Ambulatory Care Clinics

A 12-lead ECG is a routine diagnostic test performed in almost any doctor's office or ambulatory care facility. It may be performed as part of a general, routine, or pre-operative examination. This routine ECG provides a baseline tracing to be used for comparison if problems arise with a patient. The physician or trained expert looks for changes in a tracing that may indicate different types of health problems. Table 1-1 provides a list of conditions that may be diagnosed by an ECG.

TABLE 1-1 Conditions Evaluated by the ECG

- · Disorders in heart rate or rhythm and the conduction system.
- · Presence of electrolyte imbalance.
- · Condition of the heart prior to defibrillation.
- Damage assessment during and after a myocardial infarction (heart attack).
- · Symptoms related to cardiovascular disorders, including weakness, chest pain, or shortness of breath.
- · Diagnosis of certain drug toxicities.
- Diagnosis of metabolic disorders such as hyper- or hypokalemia, hyper- or hypocalcemia, hyper- or hypothyroidism, acidosis, and alkalosis.
- · Heart condition prior to surgery for individuals at risk for undiagnosed or asymptomatic heart disease.
- Damage assessment following blunt or penetrating chest trauma or changes after trauma or injury to the brain or spinal cord.
- Assessment of the effects of cardiotoxic or antiarrhythmic therapy.
- Suspicion of congenital heart disease.
- · Evaluation of pacemaker function.

Another type of ECG that may be performed in an outpatient setting is treadmill stress testing (Figure 1-3). The treadmill stress test, also known as *exercise electrocardiography*, is done to determine whether the heart gets adequate blood flow during stress or exercise. The stress test is discussed in more detail in the chapter *Cardiac Stress Testing*.

A Holter monitor is one type of ambulatory monitor. It is a small box that is strapped to a patient's waist, neck, or shoulder to monitor the heart for 24 to 48 hours while the patient is at home (Figure 1-4). Other types of ambulatory monitors, such as event monitors, may be worn up to 30 days as the patient performs normal daily activities. An event monitor is a battery-powered

Figure 1-3 This patient is performing a treadmill stress test, also known as exercise electrocardiography. During the exercise, the patient's heart and blood pressure are monitored carefully.






Figure 1-4 The Holter, a type of ambulatory monitor, allows the patient to participate in routine daily activities while the electrical activity of the heart is being recorded.

©David Page/AGE Fotostock



portable device that the patient controls to record the heart's electrical activity (ECG) when symptoms occur. Holter monitor is continued monitoring. After the monitoring period, the ECG tracing is analyzed and interpreted by the physician. Various types of ambulatory monitors are discussed in detail in the chapter *Ambulatory Monitoring*.

Outside of a Healthcare Facility

Outside of a healthcare facility, the ECG is used during cardiac emergencies such as a myocardial infarction. Emergency medical technicians and paramedics are equipped with portable ECG machines that can produce an ECG tracing at the site of the emergency. Whether the patient is at home, in a car, or in a crowded football stadium, emergency personnel can trace and monitor the electrical activity of the heart. Figure 1-5 shows one example of a portable ECG machine. In an emergency setting, the tracing can be evaluated for an abnormal ECG pattern. It is either transmitted back to the physician for evaluation or assessed by the emergency medical personnel at the scene. An abnormal pattern may require immediate treatment.

Defibrillators

Treatment for abnormal rhythms may include use of a defibrillator and/or administration of cardiac medications. A **defibrillator** produces an electrical shock to the heart that is intended to correct the heart's electrical pattern. A defibrillator is commonly used in emergencies such as a Code Blue in the hospital or other care facilities or at the site of the emergency by specially trained personnel. In situations when defibrillation is the appropriate treatment (pulseless ventricular tachycardia and ventricular fibrillation), the heart must be defibrillated quickly. The survival rate of the victim decreases by 7% to 10% for every minute a normal heartbeat is not restored.

Automatic External Defibrillators

Automatic external defibrillators (AEDs) have enabled lay rescuers to help patients with sudden cardiac arrest and serious dysrhythmias (Figure 1-6). AEDs are available in public and private places where large numbers of people gather or live. They may also be kept in homes by people who

defibrillator A machine that produces and sends an electrical shock to the heart in an attempt to correct the electrical pattern of the heart.

automatic external defibrillator (AED) A light-weight, portable device that recognizes abnormal rhythm and determines if the rhythm is considered a "shockable rhythm."

dysrhythmia Abnormal heartbeat.

Copyright © 2019 by McGraw-Hill Education

Figure 1-5 A portable ECG monitor is transported to the scene during a cardiac emergency and is attached to the patient. The ECG tracing is recorded and viewed by the emergency personnel. In addition, the tracing can be transmitted to the hospital, where a physician can evaluate and determine the necessary drugs and treatment for the patient based upon the heart rhythm viewed and the report from the emergency personnel.

©Chris Bjornberg/Science Source




Figure 1-6 Automatic external defibrillators (AEDs) can deliver an electrical impulse that may correct an abnormal heart rhythm and increase the survival rate of myocardial infarction victims. AEDs can be found in public places and require minimal training to operate.

©Ron Thompson/Rapport Press/ Newscom



are at high risk for heart attacks. An AED is a lightweight, portable device that recognizes an abnormal rhythm and determines if the rhythm is considered a "shockable rhythm." *Note:* The equipment is placed only on patients who are unresponsive to stimulation (who cannot be aroused) and have no evidence of breathing or a pulse. AEDs shock abnormal heart rhythms such as pulseless ventricular tachycardia and ventricular fibrillation. Learning about normal and abnormal rhythms is part of rhythm strip interpretation, which

cardiopulmonary
resuscitation (CPR) The
provision of ventilations
(breaths) and chest compressions (blood circulation) for a
person who shows no signs
of breathing or having a
heartbeat.

telemedicine A monitoring system in which ECG tracings are communicated from a patient outside a medical facility to the physician via a telephone or digital system.

electrocardiograph (ECG) technician An individual who has the technical skills and knowledge to record an ECG and prepare it for the physician.

is discussed in later chapters. When the machine recognizes other rhythms that cause the patient to be unresponsive, the AED recommends beginning **cardiopulmonary resuscitation (CPR).** Individuals using an AED should consider safety for themselves and the patient. A healthcare-provider-level CPR course is best for learning this technique. The patient should be checked for medication patches, pacemakers, and metal objects that could cause burns. In addition, do not use an AED when the patient is in water.

Once the equipment is placed on the patient's bare chest, it analyzes the rhythm to determine if it is likely to respond to an electric shock. Once the machine has positively identified the abnormal rhythm, it may indicate that a "SHOCK IS ADVISED." Everyone near the patient must move back and not touch the patient. One person will then announce, "clear," "everyone clear," and press the shock button. After the shock has been provided, the rescuers continue administering CPR until the patient wakes up, the machine indicates to defibrillate again, or specially trained healthcare professionals take over. AEDs make it possible for laypeople to perform defibrillation safely. The AED is now viewed as a necessary piece of equipment—similar to a fire extinguisher.

Telemedicine

Another use of the ECG tracing outside of a healthcare facility is **telemedicine**. In telemedicine, ECG tracings are communicated to the physician via the telephone or digital system. *Transtelephonic monitoring* means transmitted (trans) over the telephone (telephonic). The improvements in solid-state digital technology have expanded transtelephonic transmission of ECG data and enhanced the accuracy of software-based analysis systems. *Digital monitoring* allows ECG data to be recorded with an electronic device and then transmitted over the Internet to the healthcare facility. Transtelephonic monitoring requires a licensed practitioner to read and evaluate the tracing, whereas the digital monitoring provides a report that is validated by the licensed practitioner.

Both of these types of monitoring help physicians evaluate the ECG tracing of a patient over time. They are useful for patients with symptoms of heart disease that did not occur while they were in the healthcare facility. The recorded monitoring can be accomplished with magnetic tape (transtelephonic) or digital (computerized) recordings that can be used for up to 30 days. A transtelephonic monitor is connected to a digital phone, and the ECG is transmitted to a healthcare facility on specific days throughout the monitoring period (Figure 1-7). Individuals using a transtelephonic monitor must understand when and how to record and send a transmission.

Opportunities in Electrocardiography

Many healthcare professionals work with electrocardiography as a part of their profession. Some examples include medical assistants, nurses, emergency medical technicians, and paramedics. There are a few careers in which people work exclusively with the ECG. These include the ECG technician, the ECG monitoring (telemetry) technician, and the cardiovascular technologist.

An **electrocardiograph (ECG) technician** is an individual who records the ECG and prepares the report for the physician. ECG technicians should be able to determine if a tracing is accurate and recognize abnormalities caused

Figure 1-7 Transtelephonic monitoring uses a cellular phone device (circled) to transmit the patient's ECG tracing to a central location for monitoring.

©Total Care Programming, Inc.



cardiovascular
technologist An individual
who has advanced skills and
can assist physicians with
invasive cardiovascular diagnostic tests and procedures,
such as angioplasty or heart
surgery.

healthcare professional to

treat abnormalities.



by interference during the recording procedure. Most ECG technicians are employed in hospitals, but they may also work in medical offices, cardiac centers, cardiac rehabilitation centers, and other healthcare facilities. In some large hospitals, ECG technicians work in the home healthcare branch. They take the ECG machine to the patient's home; record the ECG; and give or transmit the report to the physician for interpretation. With the development of multiple tests to evaluate the heart, the ECG technician who obtains continuing education can expect a rewarding career.

ECG monitor (telemetry) technicians view, analyze, and report the electrical activity of multiple patients' hearts on a monitor (Figure 1-8). ECG monitor technicians are employed at hospitals or other inpatient facilities where patients are attached to continuous or telemetry monitors. The main responsibility of an ECG monitor technician is to view the ECG tracings and, if an abnormal heart rhythm occurs, alert the healthcare professional who can treat the abnormality. ECG monitor technicians must be able to evaluate the ECG tracing. They must understand the various heart rhythms and recognize abnormal ones. They may also be asked to perform other duties, such as maintaining patient records and recording ECGs.

If you enjoy the field of electrocardiology and want to advance your skills or education, you may choose to be a **cardiovascular technologist**. Technologists require more extensive training than technicians. They may assist physicians with invasive cardiovascular diagnostic tests and procedures such as angioplasty, heart surgery, or implantation of electronic, artificial pacemakers. Another specialization for cardiovascular technologists is performing ultrasounds on the heart and/or blood vessels. Ultrasound equipment transmits sound waves and then collects the echoes to form an image on a screen. As part of their duties, cardiovascular technologists may also perform ECGs.

Figure 1-8 Multiple patients can be monitored on a single monitor screen. The patients being monitored may be in a hospital or at home depending upon the type of monitoring device they are using.

©Paul Burns/ Corbis RF





- **1.** An automatic external defibrillator (AED) is used to treat what conditions?
- **2.** Briefly describe the role of an ECG technician.

1.3 **Preparing for an ECG**

Preparing for an ECG involves more than just making sure the equipment is in order. You must be aware of legal and ethical issues, and you must also communicate appropriately with the patient to make sure the patient understands the procedure.

Legal and Ethical Issues

Laws are rules of conduct that are enforced by a controlling authority such as the government. An unlawful act can result in loss of your job, a fine, or other penalty such as time in jail. Ethics are standards of behavior and concepts of right and wrong. They are based on moral values that are formed through the influence of the family, culture, and society. Unethical acts may result in poor

law Rules of conduct enforced by a controlling authority such as the government.

ethics Standards of behavior and concepts of right and wrong.

job evaluations or even job loss. When comparing law and ethics, you should understand that illegal acts are always unethical, but unethical acts are not always illegal.

Protecting Patient Information: HIPAA

In 1996, the Health Insurance Portability and Accountability Act (HIPAA) was established in response to patient information that was being transferred electronically for medical transactions. This act established a national standard for electronic healthcare transactions and also for providers, health plans, and employers. Its purpose was to ensure that the widespread use of electronic data was limited and secured. The patient can specify who can see information and what information is protected. A patient's information cannot be shared among healthcare professionals unless it is necessary for the patient's treatment.



Keep Information Private

The patient's chart or a computer screen with patient data should not be left out or open in an area where other patients or visitors may be able to view it. This is a breach of confidentiality and HIPAA.

Practicing Ethics

Many professions have a *code of ethics*. These are standards of behavior or conduct as defined by the professional group. As a healthcare professional, you must follow the standards of behavior or code of ethics set forth by your profession and place of employment. The following are some basic ethics you should practice.

Confidentiality is an essential part of patient care. You may collect information about a patient for use during his or her care and treatment; however, this information should not be made public. Confidentiality is a basic right of every patient. You should not speak about your patient or allow information about your patient to be heard or seen by anyone other than those caring for him or her. A breach in confidentiality is both unethical and illegal.

Treat all patients with respect and dignity. You should respect the privacy of patients at all times. Avoid exposing your patient's body when performing any procedure by closing the door, pulling the curtain, and/or draping the patient. In some cases, it may be necessary for a male healthcare professional to have a third person present when performing an ECG on a female. Check the policies at the facility where you are employed. Practicing ethics also includes acting professionally and cooperating with co-workers, supervisors, and other healthcare professionals. Maintain your professionalism by continuing your education and training to provide the highest level of care for your patients.

Professional Liability

Medical professional liability means that a healthcare professional is legally responsible for his or her performance. Healthcare professionals can be held accountable for performing unlawful acts (malfeasance), performing

medical professional liability Legal responsibility of healthcare professionals

for their performance.

Copyright © 2019 by McGraw-Hill Education

legal acts improperly (*misfeasance*), or simply failing to perform an act (*nonfeasance*) when they should. For example, if you find a patient's wallet after he or she leaves and you decide to keep it, this is an illegal act. While you are assisting with a treadmill stress test, if you report the blood pressure results incorrectly, resulting in the patient having a severe heart attack, this is performing a legal act improperly. If you decide to take a break when you are supposed to be monitoring a patient's heart rhythm and during the time you are gone the patient experiences an abnormal heart rhythm resulting in death from lack of prompt treatment, you have failed to perform your duties as required. Always work within your scope of practice and the standards set by your profession.

Slander and Libel

You will be speaking and writing about patients as part of your job as an electrocardiographer; you should never speak defamatory words about patients even when they upset you. Making derogatory remarks about a patient—or anyone else—that jeopardizes his or her reputation or means of livelihood is called **slander**. Slander is an illegal and unethical act that could cause you to lose your job. If you write defamatory words, this is known as **libel**, which is also illegal and unethical.

Documentation

Medical care and treatment must be documented as part of the medical record. The medical record is for communication about a patient but can be used in court as evidence in a medical professional liability case. To protect yourself legally and to provide continuity of patient care, be sure to include complete information in the medical record. Each entry must be clear, accurate, legible, dated, and signed. Table 1-2 contains a list of the information that needs to be documented in the medical record.

Medical records may be paper or electronic. Electronic records are digitally signed. Electronic records improve communication and the quality and efficiency of healthcare. For example, an ECG recorded electronically can be transferred through a wireless connection immediately to a healthcare provider on the other side of the country.

slander Making derogatory remarks about someone that jeopardizes his or her reputation or means of livelihood; slander is both illegal and unethical.

libel Writing defamatory words about someone; this is both illegal and unethical.

TABLE 1-2 Required Entries for Medical Records

- Patient identification, including full name, medical record number or Social Security number, birth date, full address and telephone number, marital status, and place of employment, if applicable.
- · Patient's medical history.
- · Dates and times of all appointments, admissions, discharges, and diagnostic tests (such as an ECG).
- · Diagnostic test results.
- · Information regarding symptoms and reasons for appointment, diagnostic tests, and admissions.
- · Physician examinations and records of results, including patient instructions.
- · Medications and prescriptions given, including refills.
- · Documentation of informed consent when required.
- Name of legal guardian or representative, if patient is unable to give informed consent.

Consent

Before you can perform an ECG on any patient, the patient must agree, or consent, to having the procedure done. Consent is often implied between the patient and healthcare professional. For example, when a patient comes to the physician's office, he or she is agreeing to be treated by the physician. This is implied consent. When a patient agrees to the ECG procedure, this is also implied consent.

Think It Through



Obtaining Consent

When a patient who cannot read is required to sign a consent form, you will need to explain the procedure to both the patient and a family member or the patient's legal guardian or representative. The patient then signs the consent form. If the patient cannot write, explain the procedure to the patient with a witness present. Ask the patient to place an *X* on the form in the witness's presence, and then have the witness sign.

Who should sign the consent form if a patient cannot read or write?

Medical procedures such as surgery and certain diagnostic tests, including a treadmill stress test, require *informed consent*. The patient must understand the procedure, benefits, alternative procedures, associated risks, and potential risk if there is a refusal of treatment. Informed consent requires the patient to sign a consent form.

Patient Education and Communication

Communicating with your patients is key to recording an ECG successfully. You must develop a positive relationship and atmosphere to reduce apprehension and anxiety during an ECG. You can reduce the patient's fears and make the ECG a positive experience by practicing effective communication techniques. Maintain a friendly, confident manner while interacting with your patient.

Helping the patient understand the procedure and follow instructions is essential. When explaining the procedure, use simple terms and speak slowly and distinctly. Encourage the patient to ask questions and repeat the instructions. This process will help ensure patient understanding. Your patient will be more cooperative if he or she trusts that you are competent to perform your job.

Communicate & Connect



Improving Communication

When speaking to a patient who is hard of hearing, look directly at the patient and speak slowly and distinctly. The patient may be able to read your lips. If the patient speaks another language, you may want to ask an interpreter or family member to assist you with communication, thus reducing apprehension and anxiety.

Think It Through



Critical Thinking and Problem Solving

Being able to troubleshoot situations that arise during the ECG procedure is essential. Troubleshooting requires critical thinking. Critical thinking is the process of thinking through the situation or problem and making a decision to solve it. The problem-solving process includes the following steps:

- 1. Identify and define the problem.
- 2. Identify possible solutions.
- 3. Select the best solution.
- **4.** Implement the selected solution.
- **5.** Evaluate the results.
- **6.** If, in step 5, you determine that the problem has not been solved, repeat steps 2 through 5 until an acceptable solution is reached.

While performing an ECG, you may need to troubleshoot actual or potential complications using the steps of the problem-solving process. These problems may arise from the patient's condition, lack of patient communication, equipment failure, or other complications. For example, suppose that you are about to perform an ECG, and the patient refuses to let you attach the lead wires. As part of troubleshooting, you ask the patient why she is refusing. The patient states, "I do not want that electricity going through me!" In a calm manner, you explain that the machine does not produce or generate electricity, and it is not harmful. After your explanation, the patient agrees to have the ECG. You have performed successful troubleshooting.

The previous example concerns a problem with communication. However, you may also need to troubleshoot problems that occur with the equipment or tracing produced. Throughout this text, the Think It Through boxes will provide a variety of problems or situations you may encounter and then ask for your solution. Use your critical thinking and problemsolving skills to answer each question. In each chapter, review the "What Should You Do?" questions to check your ability to think critically and troubleshoot.

Checkpoint
Question
(LO 1.3)

1. Explain how you would employ the steps of the problem-solving process if a patient refuses to have an ECG.

Copyright © 2019 by McGraw-Hill Education

Copyright © 2019 by McGraw-Hill Education

1.4 Infection Control

Healthcare-associated infections (HAIs) are infections that occur while patients are receiving medical or surgical care. HAIs were previously known as nosocomial infections. HAIs are a threat to patient safety. Preventing the spread of infection is an essential part of providing healthcare and performing an ECG. The Centers for Disease Control and Prevention (CDC) has implemented two levels of precautions to prevent infections—standard precautions and isolation precautions.

Standard Precautions

Standard precautions include a combination of performing hand hygiene and wearing gloves when there is a possibility of exposure to blood and body fluids, nonintact skin, or mucous membranes (Figure 1-9). Standard precautions apply to blood and all body fluids, secretions, and excretions (except sweat), regardless of whether they contain visible blood, and are practiced in all employment situations in which exposure to blood or body fluids is likely. Standard precautions reduce the risk of transmission of microorganisms from both recognized and unrecognized sources of infection. In addition to performing hand hygiene and wearing gloves, practices may include using **personal protective equipment (PPE)** such as a gown, mask, and eye protection (Figure 1-10).

In addition, the CDC advises healthcare workers not to wear artificial nails because they are more likely to harbor gram-negative pathogens than natural nails, both before and after handwashing. Natural nails should be no more than 1/4-inch long.

Table 1-3 provides a list of standard precautions that you should practice when recording an ECG. See the appendix *Standard and Isolation Precautions* for additional information about these precautions. As a rule, it is better to wear gloves even if it is not necessary. Do not forget to wash your hands or perform hand hygiene before putting on gloves and after removing them.

standard precautions

Procedures, used with all patients, that are designed to prevent the spread of infection, such as performing hand hygiene and wearing gloves.

personal protective
equipment (PPE) Devices
such as gloves, gowns,
face masks or shields, and
eye protection designed to
protect a healthcare worker
from sources of infection.

Figure 1-9 A. Handwashing, especially between patients, is essential to prevent the spread of infection. B. The use of an alcohol-based rub on hands without visible soilage is an accepted technique for preventing infection.

A. ©McGraw-Hill Education/Jill Braaten, photographer, B. ©Lillian Mundt





B.

Copyright © 2019 by McGraw-Hill Education

Figure 1-10 Wearing appropriate personal protective equipment (PPE) reduces the risk of transmission of infection. PPE includes items such as gloves, mask, gown, and eye protection.

©Total Care Programming, Inc.




TABLE 1-3 Standard Precautions Related to Electrocardiography

Hand Hygiene

- · Wash your hands after touching blood, body fluids, secretions, excretions, and contaminated items.
- Wash your hands before putting on gloves and after removing gloves.
- · Wash your hands between patient contacts.
- Wash your hands between tasks and procedures on the same person.
- Use alcohol-based hand rub if you have no visible soilage.

Gloves

- Wear gloves when touching blood, body fluids, secretions, excretions, and contaminated items.
- · Wear gloves when touching mucous membranes and nonintact skin.
- Change gloves between procedures and patients.
- Change gloves after contacting materials that are highly contaminated.
- · Remove gloves promptly after use.
- Remove gloves before touching uncontaminated surfaces or items.
- · Wash your hands immediately after glove removal.

Masks, Eye Protection, and Face Shields

• Wear mask, eye protection, and face shield during procedures and tasks that are likely to cause splashes or sprays of blood, body fluids, secretions, and excretions.

(Continued)

(Continued)

Gowns

- Wear a gown during procedures and activities that are likely to cause splashes or sprays of blood, body fluids, secretions, or excretions.
- · Remove a soiled gown promptly.
- · Wash your hands immediately after gown removal.

Equipment

- · Handle used equipment carefully. It may be soiled with blood, body fluids, secretions, and excretions.
- Prevent skin and mucous membrane exposure and clothing contamination.
- · Clean, disinfect, or sterilize reusable equipment before it is used on another person.
- · Discard single-use equipment promptly.

Environmental Control

• Follow facility procedures for the routine care, cleaning, and disinfection of surfaces. This includes environmental surfaces, nonmovable equipment, and other frequently touched surfaces.

Occupational Health and Bloodborne Pathogens

· Use resuscitation devices for mouth-to-barrier resuscitation.

Safety & Infection Control



Hand Hygiene

Performing proper hand hygiene is the single most important thing you can do to prevent the spread of infection. Wash your hands or use an alcoholbased rub (if no visible soilage is present) between patients and procedures and before and after you use gloves. *Note:* Certain types of infections, such as *Clostridium difficile*, require handwashing because the use of alcoholbased hand rubs is not sufficient to kill all the infectious organisms. Always use the method of hand hygiene that is most appropriate for the patient's condition.

isolation precautions

The second level of steps taken to prevent the spread of infection; some examples include separating the infected patient from others and using personal protective equipment.

Isolation Precautions

Isolation precautions make up the second level of protection. These precautions are based on how the infectious agent is transmitted. Isolation precautions include the following:

- Airborne precautions—require special air handling, ventilation, and additional respiratory protection (HEPA or N95 respirators).
- Droplet precautions—require mucous membrane protection (goggles and masks).
- Contact precautions—require gloves and gowns for direct skin-to-skin contact or for contact with contaminated linen, equipment, and so on.

Isolation precautions are used less often than standard precautions and only with patients who have specific infections. When isolation precautions are in place for a patient during an ECG, you will be required to follow the specific guidelines for the type of precautions implemented. Follow the facility's policy and the guidelines provided in Table 1-4.

Airborne Precautions

For patients known or suspected to be infected with microorganisms transmitted by minute airborne droplets:

- Use a private room that has monitored negative air.
- Keep the room door closed and the patient in the room.
- Wear special HEPA N-95, N-99, or N-100 masks for respiratory protection when entering the room of a patient with known or suspected infectious pulmonary tuberculosis.
- Do not enter the room of a patient known or suspected to have rubeola or varicella if you are susceptible; if you must enter the room, wear respiratory protection.
- · Limit the movement and transport of patients from the room to essential purposes only.
- Place a surgical mask on the patient if transport or movement is necessary.

Droplet Precautions


For patients known or suspected to be infected with microorganisms transmitted by droplets that can be generated by the patient during coughing, sneezing, talking, or the performance of procedures.

- · Place patient in a private room (special air handling and ventilation are not necessary and the door may remain open).
- Wear a mask when working within 3 feet of the patient.
- · Limit the movement and transport of the patient from the room to essential purposes only.
- Use a mask on the patient if transport or movement is necessary.

Contact Precautions

Contact precautions are used for patients known or suspected to be infected or colonized with microorganisms that can be transmitted by direct or indirect contact. *Direct contact* includes hand or skin-to-skin contact that occurs when performing patient care that requires touching the patient's dry skin; *indirect contact* includes touching environmental surfaces or patient-care items in the patient's environment. For these patients:

- · Place patients in a private room.
- · Wear gloves according to standard precautions.
- Wear gloves when entering the room and while providing patient care.
- Change gloves after having contact with infective material that may contain high concentrations of microorganisms, such as feces and wound drainage.
- Remove gloves before leaving the patient's room.
- · Wash hands immediately.
- · Do not touch potentially contaminated environmental surfaces or items in the patient's room after glove removal.
- All equipment must be disinfected after leaving the patient's room. The 12-lead ECG machine must be wiped down immediately.



Copyright © 2019 by McGraw-Hill Education

vital signs Temperature, pulse, respirations, blood pressure, and pain assessment.

cardiac output The volume of blood the heart pumps each minute.

1.5 Vital Signs

A patient's **vital signs**—pulse, respiration, blood pressure, temperature, pulse oximetry, weight, and pain assessment—are among the most important assessments for determining a patient's current health status. Changes in the vital signs can indicate an abnormality, or they can be a normal response to exertion, heat, stress, or other environmental factors. This section focuses on adult pulse, respirations, and blood pressure because these vital signs may be checked or monitored by ECG professionals. Table 1-5 shows the normal ranges for vital signs in adults.

Pulse and Respiration

Pulse and respiration are related because the circulatory and respiratory systems work together. Pulse is measured as the number of times the heart beats in 1 minute. Respiration is the number of times a patient breathes in 1 minute. One breath, or respiration, equals one inhalation and one exhalation. Usually, if either the pulse or respiration rate is high or low, the other is also.

Pulse

A pulse rate gives information about the patient's cardiovascular system. It is an indirect measurement of the patient's **cardiac output**, or the amount of blood the heart is able to pump in 1 minute. If the pulse is weak, irregular, or abnormally fast or slow, the patient may show signs of low cardiac output.

Measure the pulse of adults at the radial artery, where it can be felt in the groove on the thumb side of the inner wrist (Figure 1-11). Press lightly on this pulse point with the pads of your fingers. Do not use your thumb because your thumb has a pulse, which you may feel instead of the patient's pulse. Count the number of beats you feel in 1 minute and note the rhythm and volume. The rhythm can be regular or irregular. The volume can be weak, strong, or bounding. A bounding pulse feels like it is leaping out and then quickly disappearing with each pulse beat.

In some instances you will not be able to feel a pulse at the wrist, especially if the patient is very weak or has unstable vital signs. In these cases, you can check for a pulse at the carotid artery. This artery is located on the side of the neck next to the airway (Figure 1-12). This is the same pulse location that is used for adults during basic life support.

TABLE 1-5 Vital Signs in Adults

Pulse: 60 to 100 beats per minute	
Respirations: 12 to 20 breaths per minute	
Blood pressure (mm Hg)	
Systolic	Diastolic
Less than 120 (normal)	Less than 80 (normal)
120 to 139 (prehypertension)	80 to 89 (prehypertension)
140 to 159 (Stage 1 hypertension)	90 to 99 (Stage 1 hypertension)
160 or higher (Stage 2 hypertension)	100 or higher (Stage 2 hypertension)

Figure 1-11 Place your fingers in the groove on the thumb side of the inner wrist to check a patient's pulse.





Figure 1-12 To check the carotid pulse, place your fingers on the side of the neck next to the airway.



Respiration

A patient's respiration rate indicates how well the patient's body is providing oxygen to tissues. The best way to check respiration is by watching, listening, or feeling the movement at the patient's chest, stomach, back, or shoulders. If you cannot see the chest movement, then place your hand over the patient's chest, shoulder, or abdomen and listen and feel for the movement of air. Respirations also may be counted with a stethoscope. Place the stethoscope on one side of the spine in the middle of the back to count respirations.

You need to count respirations subtly because once the patient is aware that respiration is being measured, he may unintentionally alter his breathing. If you are using a stethoscope, tell the patient that you want to listen to his lungs. If you are not using a stethoscope, count the respirations while you have your hand on the pulse site.

Respirations are counted for 1 full minute to determine the rate, rhythm, and effort (quality). Counting for less than a minute may cause you to miss certain breathing abnormalities. The rhythm should be regular. The quality or effort may be normal, shallow, or deep. Irregularities such as *hyperventilation* (excessive rate and depth of breathing), *dyspnea* (difficult or painful breathing), *tachypnea* (rapid breathing), or *hyperpnea* (abnormally rapid, deep, or labored breathing) are indications of possible disease and should be noted.

Rales, rhonchi, and gurgling are types of noisy breathing that can indicate an abnormality. Rales—crackling sounds—may indicate fluid in the

systolic blood pressure

The blood pressure measured when the left ventricle of the heart contracts; the number corresponding with the first Korotkoff sound heard when performing an auscultated blood pressure.

diastolic blood pressure

The blood pressure measured when the heart relaxes; the minimum amount of pressure exerted against the vessel walls at all times.

hypertension High blood pressure.

hypotension A lower than normal blood pressure that can cause reduction of blood flow to vital organs.

tilt table test A test to determine whether a change in a patient's blood pressure or heart rate due to a change of position is causing symptoms of lightheadedness or fainting. lungs and can be heard in patients with pneumonia, congestive heart failure, and other conditions. Rhonchi are wheezing or snore-like sounds that occur when the airways are narrowed or obstructed.

Blood Pressure

Blood pressure (also known as *arterial blood pressure*) is the force at which blood is pumped against the walls of the arteries. The standard unit for measuring blood pressure is millimeters of mercury (mm Hg). The pressure measured when the left ventricle of the heart contracts is known as the **systolic blood pressure**. The pressure measured when the heart relaxes is known as the **diastolic blood pressure**. The diastolic pressure indicates the minimum amount of pressure exerted against the vessel walls at all times.

According to the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure, expected adult systolic readings are less than 120 mm Hg and adult diastolic readings are less than 80 mm Hg. These values may increase with advancing age. A systolic reading between 120 and 139 mm Hg and a diastolic reading between 80 and 90 mm Hg is considered prehypertension. A reading between 140 mm Hg and 159 mm Hg systolic and between 90 mm Hg and 99 mm Hg diastolic is considered Stage 1 hypertension. A systolic reading above 160 mm Hg and above 100 mm Hg diastolic is considered Stage 2 hypertension. Table 1-5 outlines these blood pressure classifications.

Factors Affecting Blood Pressure

Many factors can affect blood pressure. Risk factors such as being overweight or obese, lack of physical activity, too much salt in the diet, too much alcohol consumption (more than one to two drinks a day), stress, age, and genetics can play a role in increasing the blood pressure. Internal factors such as cardiac output, blood volume (amount of blood in the body), vasoconstriction (constriction of the blood vessels), presence of atherosclerotic plaque in the blood vessels, and blood viscosity (thickness) can impact the blood pressure within the body.

If a patient's blood pressure reading is consistently elevated after two or more visits to the physician, the patient may be diagnosed with **hypertension**, or high blood pressure. Hypertension may be categorized as essential or secondary. *Essential hypertension* has no identifiable cause and is sometimes referred to as *idiopathic*. According to the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure, part of the National Heart, Lung, and Blood Institute in the National Institutes of Health, 95% of all hypertension is essential. *Secondary hypertension* occurs as a result of some other condition, such as kidney or heart disease.

Hypotension, or low blood pressure, is not generally a chronic health problem. Slightly low blood pressure may be normal for some patients and does not usually require treatment. Severe hypotension may be present with dehydration, shock, heart failure, severe burns, and excessive bleeding.

People who have severe hypotension, a very slow heart rate, or both may have frequent fainting episodes. For patients complaining of feeling faint or lightheaded, the practitioner may order a **tilt table test**. In this test, the patient lies quietly on a table, and straps are applied to hold the patient in

auscultated blood pressure Blood pressure determined while listening with a stethoscope. place. After a short time, the table is suddenly tilted upright to simulate moving from a lying position to standing, and the table remains at that angle until the patient's heart rate and blood pressure stabilize. The patient's heart rate and blood pressure are monitored during the entire test to determine whether a drop in heart rate or blood pressure is causing the person's symptoms.

Measuring Blood Pressure

Blood pressure is measured using a *sphygmomanometer*, or blood pressure cuff. To measure blood pressure, wrap the blood pressure cuff around the patient's upper arm, 1 to 2 inches above the pulse point of the brachial artery (Figure 1-13). This pulse point is located in the bend of the elbow, called the *antecubital space*. Make sure the cuff is lined up over the brachial artery to ensure the inflation bladder is centered for an accurate blood pressure reading. Never apply the blood pressure cuff over clothing or place the stethoscope under the blood pressure cuff. Ensure that the patient's legs are not crossed and the feet are flat on the floor during the measurement.

When measuring blood pressure, you may need to determine the palpatory pressure that represents the target peak inflation. Palpation provides an approximation of the systolic blood pressure to ensure an adequate level of inflation when the actual measurement is made. It also helps prevent overinflation, which is uncomfortable for the patient. In most situations you may simply inflate the cuff to 180 to 200 mm Hg or at least 30 mm Hg over the highest recorded systolic blood pressure for your patient.


When palpatory pressure is required, inflate the cuff as you palpate the radial pulse. Inflate the cuff until you can no longer feel the radial pulse and note the pressure at that point. Release the air from the cuff and allow the arm to rest for 30 to 60 seconds, or remove the cuff entirely and replace it after 30 to 60 seconds.

To determine the **auscultated blood pressure**, inflate the cuff to 30 mm Hg above the palpatory result, generally in the range of 180 to 200 mm Hg. Then, while you slowly release the air in the cuff (2 to 4 mm Hg/second), listen with the stethoscope placed over the brachial pulse point. You will hear vascular sounds (thumping) that will change. These sounds are called *Korotkoff sounds*. Record the number when you hear the first sound. This is the systolic number. Record the number when you hear the final sound. This is the diastolic number.

Figure 1-13 A. Place the blood pressure cuff about 2 inches above the bend in the patient's elbow. B. Palpate the blood pressure at the antecubital space.

©Jeffrey Coolidge/Getty Images RF





В

Checkpoint
Questions
(LO 1.5)

1. Which three vital signs are of particular importance to ECG technicians?

2. What is the difference between a palpated blood pressure and an auscultated blood pressure?

Chapter Summary

Learning Outcomes	Summary	Pages
1.1 Describe the history and the importance of the ECG.	An ECG is a tracing of the heart's electrical activity made using an electrocardiograph. Willem Einthoven is credited with developing the first electrocardiograph. Today, the machine is used to diagnose cardiovascular disease, which is the number one cause of death in America.	2–3
1.2 Identify the uses of an ECG and opportunities for an electrocardiographer.	The ECG is used routinely in healthcare facilities before surgery, during emergencies, and as part of continuous monitoring. In an outpatient setting, an ECG is part of an exam or is used during stress tests or Holter monitoring. Outside of the healthcare facility, the ECG is used to evaluate patients cardiac status and during emergencies. Career opportunities in electrocardiography include ECG technician, ECG monitor technician, and cardiovascular technologist.	3–10
1.3 Troubleshoot legal, ethical, patient education, and communication issues related to the ECG.	Legal and ethical issues include protecting patient information in accordance with HIPAA, preventing situations that lead to professional liability, avoiding slander and libel, documenting all procedures thoroughly, and obtaining the patient's consent before performing any procedure. Communicating with and educating the patient are an important part of preparing for an ECG. Troubleshooting may involve solving problems with the patient, the equipment, or the tracing produced.	10–14
1.4 Perform infection control measures required for the ECG.	Follow all infection control guidelines at your facility, including standard precautions and isolation precautions as necessary. Performing proper hand hygiene is the single most important thing you can do to prevent the spread of infection.	15–18
1.5 Compare basic vital sign measurements related to the ECG.	The vital signs include pulse, respiration, blood pressure, temperature, pulse oximetry, weight, and pain assessment. ECG technicians are commonly asked to check or monitor the pulse, respiration, and blood pressure of adults. Normal vital signs for most adults are a pulse of 60–100 beats per minute, 12–20 respirations per minute, and a blood pressure of less than 120 mm Hg (systolic) over 80 mm Hg (diastolic).	19–23

Chapter Review

Match	ning	
	1. cardiovasc disease (LC) 2. electrocard (LO 1.1) 3. coronary adisease (LC) 4. standard production (LO 1.4) 5. electrocard (LO 1.1) 6. defibrillato 7. AED (LO 1.8)	heart b. a tracing of the signal produced by the heart's electrical activity and used for diagnostic evaluation of the heart c. the second level of protection against the transmission of disease; based on how the infectious agent is transmitted d. narrowing of the heart's blood vessels that causes a reduction of blood flow to the heart e. used to analyze the heart rhythm and produce a shock if necessary f. disease related to the heart and blood vessels (veins and arteries) g. a machine that delivers an electrical shock to the heart that is intended to correct the abnormal electrical pattern of the heart
True/i	8. isolation pr (LO 1.4)	, and the second
	ch statement and provided.	determine if it is true or false. Circle the T or F. Correct false (F) statements on
т ғ	9. An ECG ma	chine records the electrical activity of the heart. (LO 1.1)
т ғ	_	ecautions are guidelines written for healthcare providers to help prevent the fection. (LO 1.4)
T F	11. Libel is the	ct of making derogatory remarks about someone. (LO 1.3)

Т	F	12.	A transtelep	ohoni	ic monitor transmits an ECG over the Internet. (LO 1.2)
Т	F	13.	Because an	ECG	is not an invasive procedure, it is not necessary to wear personal protective
) when performing this procedure under any circumstances. (LO 1.4)
M	.141.	nlo i	Choice		
			Choice	70	

Circle the correct answer.

- **14.** The normal range for an adult's pulse is (LO 1.5)
 - **a.** 40–80 beats per minute.
 - **b.** 50–60 beats per minute.
 - c. 60–100 beats per minute.
 - **d.** 90–120 beats per minute.
- 15. Which physician received a Nobel Prize for inventing the electrocardiograph? (LO 1.1)
 - a. Augustus D. Waller
 - **b.** Sir Thomas Lewis
 - c. Willem Einthoven
 - d. Joseph Lister
- **16.** The main responsibility of an ECG monitor technician is to (LO 1.2)
 - a. view and evaluate ECG tracings on an electronic display and alert healthcare professionals when abnormalities appear.
 - **b.** attach the ECG electrodes to patients who will be monitored remotely.
 - c. assist physicians with angioplasty, heart surgery, and implantation of artificial pacemakers.
 - **d.** perform ultrasounds on patients' blood vessels.
- 17. Which of the following is a reason for performing an ECG? (LO 1.2)
 - a. To evaluate heart conditions
 - **b.** To check for problems with the flow of blood through the heart
 - c. To see how well the veins are contracting
 - **d.** To evaluate the rate and rhythm of breathing
- **18.** A defibrillator can be used (LO 1.2)
 - **a.** to treat an abnormal heart rhythm.
 - **b.** without training.
 - **c.** to produce an electrical rhythm.
 - **d.** to record the patient's breathing pattern.
- **19.** Transtelephonic monitoring allows for information to be (LO 1.2)
 - **a.** reviewed immediately by the patient.
 - **b.** transmitted over a telephone.
 - c. submitted for billing purposes only.
 - **d.** recorded by a computerized device.

Education
aw-Hill
McGra
e by
201
9
opyrigh
Ö

28.	which of the following is not considered a vital sign? (LO 1.5) a. Blood pressure b. Pain assessment c. Urinary output d. Respirations
29.	 Which of the following is the most common pulse point for measuring an adult's pulse? (LO 1.5) a. Radial artery b. Popliteal artery c. Brachial artery d. Temporal artery
30.	The systolic blood pressure represents the blood pressure when (LO 1.5) a. the right atrium relaxes. b. the left ventricle contracts. c. the right ventricle relaxes. d. the left atrium contracts.
31.	A patient's pulse is an indirect measurement of his or her (LO 1.5) a. blood volume. b. blood viscosity. c. vasoconstriction. d. cardiac output.
	 Which of the following is not a specific type of isolation precaution? (LO 1.4) a. AIDS precautions b. Droplet precautions c. Airborne precautions d. Contact precautions
Crit	ical Thinking Application What Should You Do?
	the following situations and use critical thinking skills to determine how you would handle each. your answer in detail in the space provided.
33.	You have been performing ECGs at a local clinic for about 6 months. Your favorite uncle says to you, "Since I just turned 40, your Aunt Beth thinks I should have an ECG. Will you do one on me if I come by where you work?" What would you say or do for your uncle? Consider the following: (LO 1.3) Should your uncle have an ECG?
	Should you do the ECG if he stops by your office? Why or why not?

34.			ing some mild chest pain. During his ECG, he says, "How does it look? Is What would be your best response? (LO 1.3)
5.			ere a co-worker is performing an ECG on a female patient. The door is open overed. What would you do? (LO 1.3)
86.		u are	monitoring the heart rhythms on six patients at a local hospital when you in desperate need to go to the restroom, and you really want to go home. O 1.3)
W	connect		w that you have completed the material in the textbook, go to Connect and mplete any chapter activities you have not yet done.

Competency Checklists

Use the following Competency Checklists 1-1 and 1-2 to practice and perform the skills presented in this chapter. (LO 1.5)

COMPETENCY CHECKLIST 1-1

Measuring Pulse and Respirations

Yes	No	Yes	No	Yes	No	Date	Initials

COMPETENCY CHECKLIST 1-2

Measuring Adult Blood Pressure

Procedure Steps (Rationale)	Practice		Prac	tice	Te	st	Mag	stered
Preprocedure	Yes	No	Yes	No	Yes	No	Date	Initials
Gather the equipment. Check the sphygmomanometer to be sure it is calibrated and in working order. (Accurate results can be obtained only when the sphygmomanometer is correctly calibrated.)	103				163		Dute	mitals
Introduce yourself and identify the patient using two forms of identification.								
3. Wash your hands and explain the procedure to the patient.								
4. Select the appropriate size cuff for the patient. The bladder in the cuff should encircle at least 80% of the arm. If you are not sure which size cuff to use, use a larger cuff. (Using the proper cuff size increases the accuracy of the reading.)								
Procedure								
1. Wrap the cuff snugly around the patient's bare upper arm so that the midline of the bladder is over the brachial artery. The cuff should be 1 inch above the antecubital space. If the patient is wearing long sleeves, ask him or her to roll up the sleeve loosely. If the sleeve is too tight to be rolled up, ask him or her to change into a gown. (Checking the blood pressure through the clothing decreases the accuracy of the reading.)								
Position the manometer so that the gauge is at eye level and easy to see. Make sure the tubing from the blood pressure cuff is not obstructed.								
3. Close the valve on the bulb so that it is finger-tight, but do not overtighten it.								
4. Determine the palpatory pressure if required. To do so, squeeze the bulb to inflate the cuff rapidly to 70 mm Hg. Then increase the pressure in 10 mm Hg increments while using your other hand to palpate the radial pulse in the patient's wrist. Note the pressure at which the radial pulse disappears. This is the palpatory pressure. Open the valve and deflate the cuff completely. Then wait 30 seconds or remove and replace the cuff before proceeding with the auscultated blood pressure measurement. (Failure to deflate the cuff completely may cause blood to pool in the artery, which can result in an inaccurate reading.)								
5. Place the stethoscope's earpieces in your ears so that they point up or toward your nose. Rotate the head of the stethoscope in the diaphragm position and test it by tapping gently.								
6. Place the diaphragm of the stethoscope over the patient's brachial artery and hold it firmly using your index and middle fingers. Make sure the entire surface of the diaphragm is in contact with the patient's skin and that it does not touch the blood pressure cuff. (Do not use your thumb to hold the stethoscope in place because your thumb has a pulse that can interfere with the reading.)								

(Continued)

Procedure Steps (Rationale)		tice	Prac	ctice	Те	st	Mas	stered
Preprocedure	Yes	No	Yes	No	Yes	No	Date	Initials
7. Inflate the bladder rapidly to a pressure that is 20 to 30 mm Hg higher than the palpatory pressure (if obtained) or at least 30 mm Hg higher than the highest recorded systolic blood pressure for the patient. If neither are known, inflate cuff to 180 to 200 mgHg.								
 Partially open the screw on the valve to deflate the bladder at about 2 mm per second. Listen for the Korotkoff sounds. Note the pressure at which you first hear the repetitive sounds. This is the systolic pressure. 								
 Continue to deflate the cuff at the same rate, noting the pressure at which the sound becomes muffled, and then the pressure at which the sound disappears. The pressure at which the sound disappears is the diastolic pressure. 								
 Continue to deflate the cuff and listen for another 10 mm Hg. Listen to be sure the repetitive sounds do not resume. (This ensures that you are measuring the diastolic pressure, not an auscultatory gap.) 								
 Rapidly deflate the cuff the rest of the way and remove it from the patient's arm. 								
Postprocedure								
 Record the systolic and diastolic numbers in the patient's chart, separated by a slash. Record the date and time, the arm in which you measured the pressure, and the patient's position (sitting or lying). If you used a nonstandard cuff size, record the cuff size as well. 								
Use gauze moistened with alcohol to disinfect the diaphragm and earpieces of the stethoscope.								
Dispose of the used gauze appropriately and wash your hands.								
Comments:								
-								
Signed								
Evaluator:								
Student:								
Design Elements: Think It Through icon (gears in head) ©Fine Art/S (computer and stethoscope) ©lenetstan/Shutterstock.com RF; Safet								on

©Ezume Images/Shutterstock.com RF; Law & Ethics icon (gavel and stethoscope) ©Lisa S./Shutterstock.com RF.